Please check the examination details below before entering your candidate information				
Candidate surname		Other names		
Centre Number Candidate N	umber			
Pearson Edexcel Leve	l 1/Leve	el 2 GCSE (9–1)		
Monday 22 May 202	23			
Morning (Time: 1 hour 45 minutes)	Paper reference	1CH0/1F		
Chemistry		₾ •		
PAPER 1				
PAPER I				
		Foundation Tier		
You must have:		Total Marks		
Calculator, ruler		J Total Marks		

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 100.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶








BLANK PAGE

Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

- 1 In an experiment, paper chromatography was used to separate the coloured dyes in four different inks, **W**, **X**, **Y** and **Z**.
 - (a) Figure 1 shows the chromatogram at the end of the experiment.



(i) The chromatogram shows that only one of the inks contains a single dye.

Which ink contains a single dye?

(1)

- \square A W
- \square B X
- \square D Z
- (ii) Which ink contains the greatest number of dyes?

(1)

- \bowtie A W
- \boxtimes B X
- \square C Y
- \square D Z

(iii) The R_f value of a dye can be calculated using the equation

$$R_{f} = \frac{\text{distance moved by the dye}}{\text{distance moved by solvent front}}$$

At the end of the chromatography one dye had moved 3.60 cm and the solvent front had moved 9.20 cm.

Calculate the R_f value for this dye.

Give your answer to 2 decimal places.

(2)

 $R_f = \dots$

(b) The substance used as the solvent in the chromatography was heated for 8 minutes.

Figure 2 shows how the temperature of the substance changed with time.

temperature in °C




Figure 2

From **A** to **B** the substance was a liquid.

From **C** to **D** the substance was a gas.

(i) Give the name of the change when a liquid becomes a gas.

(1)

(ii) Use Figure 2 to give the temperature of the substance at 4 minutes.

(1)

(iii) Use Figure 2 to give the time when the substance has completely changed into a gas.

(1)

..... minutes

(iv) The temperature of the substance at **A** was 25 °C.

Calculate the temperature rise of the substance from **A** to **D**.

(1)

(Total for Question 1 = 8 marks)

2 (a) Most of the gold used in jewellery is not pure gold but alloys of gold.

The purity of gold is measured in carats.

Figure 3 shows how the percentage of gold is related to the purity of gold measured in carats.





Figure 3

State the relationship between the percentage of gold and the number of carats.

(1)







- This question is about electrolysis.
 - (a) Which statement describes what happens during electrolysis?

(1)

- X atoms are decomposed
- X ionic compounds are decomposed
- X mixtures are separated
- X **D** molecules are separated
- (b) Figure 5 shows the electrolysis of copper chloride solution.
 - (i) Use the words from the box to complete the labelling of the diagram in Figure 5.

(2)




Figure 5

Draw one s	straight line from each prod	uct to its name.	(2)
	product	name	
	solid X	• carbon	
		chlorine	
		copper	
	gas Y	hydrogen	
	ment is repeated using pow oride solution.	dered solid copper chloride instea	d of
Nothing ha	ppens and no products are	formed.	
Explain wh	y nothing happens and no	oroducts are formed.	(2)



4 Steel is an alloy containing iron.

When exposed to damp air, some steels will corrode to form rust.

(a) (i) Which gas in the air is needed for corrosion to occur?

(1)

- 🛚 🗛 argon
- **B** carbon dioxide
- **D** oxygen

(ii) What type of reaction happens when the iron in steel corrodes?

(1)

- A the iron has been displaced
- **B** the iron has been neutralised
- C the iron has been oxidised
- **D** the iron has been reduced

(b) Rust can be removed from steel by treating it with dilute hydrochloric acid.

One product formed in this reaction is iron chloride, FeCl₃.

Calculate the relative formula mass of this iron chloride.

(relative atomic masses: Fe = 56.0, CI = 35.5)

(2)

relative formula mass =

(c) Figure 6 shows the composition of one type of steel that has a low resistance to corrosion and another type of steel that has a high resistance to corrosion.

	percentage of element in steel		
element	steel with low resistance to corrosion	steel with high resistance to corrosion	
iron	98.2	80.6	
carbon	0.4	1.1	
chromium	0.0	17.0	
manganese	0.9	0.8	
nickel	0.5	0.5	

Figure 6

	(i)	Usina Figure	6. state which	non-metal is in	n both types	of steel.
--	-----	--------------	----------------	-----------------	--------------	-----------

(1)

(ii) Using Figure 6, state which metal is added to steel to increase its resistance to corrosion.

(1)

(iii) Explain **one** other way that corrosion of steel can be prevented.

(2)



(d) A student is given two nails of the same size but made of different types of steel. They are also given two boiling tubes and some distilled water. Devise an experiment to show which nail corrodes more quickly.	(3)
(Total for Question 4 = 11 m	arks)

- **5** (a) When lead nitrate solution and potassium chloride solution are mixed, potassium nitrate and a precipitate of lead chloride are formed.
 - (i) Complete the word equation for this reaction.

(1)

lead nitrate + _____ + lead chloride

(ii) Lead nitrate is toxic.

Which hazard symbol should be on a container of lead nitrate?

(1)











(b) A student put 5 cm³ of potassium carbonate solution into a test tube and added 2 cm³ of calcium nitrate solution.

A precipitate formed and was allowed to settle as shown in Figure 7.

The height of the precipitate was measured.




Figure 7

(i) Give the name of the piece of apparatus the student should use to find the volume of the potassium carbonate solution.

(1)

(ii) The student repeated the experiment.

The results are shown in Figure 8.

experiment	height of precipitate in cm
1	2.4
2	2.7
3	2.4

Figure 8

Use the data in Figure 8 to calculate the mean height of the precipitate.

(2)

mean height of precipitate =cm

(iii)	Describe how a pure, dry sample of the precipitate could be obtained from the mixture in the test tube.	
		(3)
(iv)	The student investigated whether increasing the volume of calcium nitrate solution increased the height of the precipitate formed.	
	They repeated the experiment using different volumes of calcium nitrate.	
	State one variable that should be controlled in this investigation.	(1)
	(Total for Question 5 = 9 ma	arks)

6	(2)	Ma	agnosium is a motal	
0	(a)		ignesium is a metal.	
		(i)	State one physical property of magnesium.	(1)
		(ii)	Which element is in the same group of the periodic table as magnesium? Use the periodic table to help you answer this question.	(1)
			A carbon	
			■ B chromium	
			C sodium	
			■ D strontium	
	(b)	(i)	Magnesium atoms have 12 electrons. Complete the electronic configuration of a magnesium atom.	(1)
		(ii)	The electronic configuration of a chlorine atom is 2.8.7	
			Explain how the electronic configuration of chlorine is linked to its period in the periodic table.	(2)
	••••••			



(c) 1.20 g of magnesium reacts completely with 3.55 g of chlorine to form magnesium chloride.	
Calculate the empirical formula of the magnesium chloride.	
(relative atomic masses: $Mg = 24.0$, $CI = 35.5$)	
You must show your working.	(3)

empirical formula =



(d) Sodium reacts with chlorine to form sodium chloride, which contains ionic bonds.

Hydrogen reacts with chlorine to form hydrogen chloride, which contains covalent bonds.

Figure 9 shows dot and cross diagrams of these compounds.

sodium chloric	de (ionic bonding)	hydrogen chloride (covalent bonding)
sodium ion	chloride ion	H CI

Figure 9

Describe the differences between an ionic bond a	
	(4)
	(Total for Operation 6 – 12 mayles)
	(Total for Question 6 = 12 marks)

7 (a) Figure 10 shows some information on a container of plant fertiliser.

contains	percentage by mass
ammonium nitrate (NH ₄ NO ₃)	46%
phosphorus oxide (P ₂ O ₅)	0%
potassium nitrate (KNO ₃)	54%

Figure 10

(i)	State which element, often present in fertilisers, is not present in this fertiliser.	

(1)

(ii) Ammonium nitrate, NH₄NO₃, is prepared for use in fertilisers by neutralising ammonia with an acid.

Which acid reacts with ammonia to produce ammonium nitrate?

(1)

- A ethanoic acid
- **B** hydrochloric acid
- C nitric acid
- **D** sulfuric acid

(iii'	State	why	farmers	spread	fertilisers	οn	their	field	ς
١	ш	Jiale	vviiy	iaiiiieis	spieau	iei uiiseis	OH	uieii	Helus	٥.

(1)

- (b) Ammonium sulfate is a fertiliser and is produced on a large scale in industry.

 In this process, ammonia reacts with sulfuric acid.
 - (i) Write the word equation for the reaction between ammonia and sulfuric acid.

(2)



(ii)	Ammonium sulfate can also be made in the laboratory by titrating
	ammonia solution with dilute sulfuric acid.

Give **one disadvantage** of using this laboratory method to produce ammonium sulfate as a fertiliser compared with an industrial method.

(1)

(6)

*(c) In the laboratory, ammonium sulfate crystals can be made using ammonia solution and dilute sulfuric acid.

The volume of ammonia solution required to neutralise 25 cm³ of dilute sulfuric acid is found by titration using an indicator.

The results of the titration can be used to prepare a solution of ammonium sulfate.

Pure, dry ammonium sulfate crystals can be made from this solution.

Figure 11 shows some of the equipment that may be used in the experiment.




Figure 11

Write a detailed method to make ammonium sulfate crystals starting with ammonia solution and dilute sulfuric acid.

 	••••												

8 In an experiment, powdered calcium hydroxide was added to dilute hydrochloric acid and the pH was measured.

The method used was

- step 1 measure 200 cm³ dilute hydrochloric acid into a beaker
- step 2 add 0.1 g of powdered calcium hydroxide to the beaker
- step 3 find the pH of the mixture
- **step 4** repeat steps 2 and 3 until the pH stops changing.
- (a) State what should be done after **step 2** to make sure that any reaction is complete.

(1)

(b) Complete the word equation for the reaction.

(2)

calcium hydroxide + hydrochloric acid →

(c) Which row of the table shows the state symbols for powdered calcium hydroxide and dilute hydrochloric acid in the balanced chemical equation?

(1)

		calcium hydroxide	hydrochloric acid
X	Α	aq	I
X	В	I	aq
X	C	S	aq
X	D	S	I





(d) The results of the experiment are shown in Figure 12.




Figure 12

(i) Using Figure 12, give the pH of the acid at the start of the experiment.

(1)

(ii) Using Figure 12, give the mass of calcium hydroxide required to make a neutral mixture.

(1)

(iii) Explain why the pH starts at a low value and ends at a higher value.

(3)





(e) State what should be used to measure the pH of the mixture in this experiment.

(1)

(f) The calcium hydroxide used is corrosive to the eyes and an irritant to skin.

Using this information, state **one** safety precaution that should be taken during the experiment when using any corrosive substance.


(1)

(Total for Question 8 = 11 marks)

BLANK PAGE



9 Figure 13 shows part of the reactivity series of metals.



increasing reactivity

Figure 13

(a) Which metal reacts when added to cold water?

(1)

- A calcium
- B copper
- C gold
- **D** silver

(b) A student investigates the reactivity of four different metals.

The student adds an equal-sized piece of each metal to separate test tubes containing dilute hydrochloric acid.

The student's observations for zinc and copper are recorded in Figure 14.

metal	observations
magnesium	
-:	bubbles produced at a steady rate
zinc	test tube feels slightly warm
iron	
copper	no reaction

Figure 14



(i)	Use the information in Figure 13 and in Figure 14 to predict the observations for the reactions of magnesium and of iron with dilute hydrochloric acid. magnesium	(2)
	iron	
(ii)	When metals react with acids, hydrogen gas is produced. Describe the test to show that the gas is hydrogen.	(2)
(iii)	When magnesium reacts with hydrochloric acid, magnesium chloride and hydrogen are formed.	
	Complete the balanced equation for the reaction.	(2)
	Mg +HCl \rightarrow MgCl ₂ +	



(6)

- *(c) There are **three** common methods of obtaining metals from the Earth's crust:
 - mine the pure metal
 - mine the metal ore and heat it with carbon
 - mine the metal ore and electrolyse the molten compound.

The method used to obtain a metal is linked to its position in the reactivity series of metals.

Aluminium, gold, iron, and silver are some commonly used metals.

Use the reactivity series in Figure 13 to state and explain the method chosen to obtain each of these four metals.

- **10** There are several stages to the production of sulfuric acid in industry.
 - (a) Sulfur dioxide is required for the production of sulfuric acid.

Sulfur dioxide can be obtained by heating copper sulfide, Cu₂S, in excess air.

$$Cu_2S + O_2 \rightarrow 2Cu + SO_2$$

Calculate the atom economy for the production of sulfur dioxide, SO₂, in this reaction.

(relative atomic mass: Cu = 63.5

relative formula masses: $O_2 = 32.0$, $Cu_2S = 159.0$, $SO_2 = 64.0$)

Give your answer to two significant figures.

(4)

atom economy =%

(b) In one stage vanadium oxide, V_2O_5 , is used.

Based on the position of vanadium in the periodic table, which row shows the most likely melting point of vanadium and colour of vanadium oxide?

(1)

		melting point of vanadium in °C	colour of vanadium oxide
X	A	50	white
X	В	1910	white
X	C	50	orange
X	D	1910	orange

TOTAL FOR PAPER = 100	MARKS
(Total for Question 10 = 11	marks)
2	
1	(2)
(iii) State two reasons why the percentage yield is less than 100%.	
percentage yield =	
Calculate the percentage yield of sulfuric acid.	(2)
The actual mass produced was 672 tonnes.	
(ii) Using a different amount of sulfur trioxide, it was calculated that 700 tonne of sulfuric acid could be made.	25
maximum mass of sulfuric acid =	tonnes
(relative formula masses: $SO_3 = 80$, $H_2SO_4 = 98$)	(2)
(i) Calculate the maximum mass of sulfuric acid that could be produced from 400 tonnes of sulfur trioxide, SO_3 .	
$SO_3 + H_2O \rightarrow H_2SO_4$	
(c) The equation shows a reaction forming sulfuric acid.	



The periodic table of the elements

_						
0	He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86
		19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85
9		16 O oxygen 8	32 S suffur 16	79 Selenium 34	128 Te tellurium 52	[209] Po polonium 84
2		14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83
4		12 C carbon 6	28 Silicon 14	73 Ge germanium 32	119 Sn th	207 Pb lead 82
က		11 B boron 5	27 Al aluminium 13	70 Ga gallium 31	115 In indium 49	204 TI thallium 81
	·			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80
				63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79
				59 Ni nickel 28	106 Pd palladium 46	195 Pt platinum 78
				59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77
	1 T hydrogen			56 iron 26	Ru ruthenium 44	190 Os osmium 76
-				55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75
		mass sol umber		52 Cr chromium 24	96 Mo molybdenum 42	184 W tungsten 74
	Key	relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73
		relativ ato atomic		48 T: titanium 22	91 Zr zirconium 40	178 Hf hafnium 72
	·			45 Sc scandium 21	89 Y yttrium 39	139 La* lanthanum 57
7		9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56
-		7 Li lithium 3	23 Na sodium 11	39 K potassium 19	85 Rb rubidium 37	133 Cs caesium 55

^{*} The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.