1	Ions are atoms that have lost or gained electrons and so become charged.	
2	Ionic compounds need to be molten or dissolved for ions to be	
2	separatedby electrolysis. An ionic substance with freely moving ions is called an electrolyte.	
3	Sand a sand a first it cory morning forto to carried all electronytes	
4	Positive ionsare called cations areattracted to	
	the negative electrodecalled the cathode .	
5	Negative ions are called anions are attracted to	
3	the positive electrodecalled the anode .	
	Half equations show what happens at the electrodes (OILRIG): ReduCtion at the Cathode, oxidAtion at Anode. Cathode reaction: Zn²² + 2e → Zn reduction—ReduCtion takes	
6	Anode reaction: 2Cl ⁻ + 2e → 2n reduction Reduction takes place at the Cathode. Note that two Cl ⁻ ions are needed to form one chlorine molecule. OxidAtion takes place at the Anode.	
7	OxidationIs theLoss of electrons andReductionIsGainof electrons during a reaction (OILRIG)	
8	Copper is purified by electrolysis, more expensive butlower resistance when used in wires	
9	Direct current (d.c.) is needed electrolysis	
10	Electrodes are washed in distilled water (remove surfaceimpurities) and dipped inpropanone(remove the water) to ensure an accurate reading on the mass balance.	

11	Molten salts undergoing electrolysis with inertelectrodes decompose into their elements. The metalions (+ve) move to the cathode (-ve), the non-metal (-ve) moves to the anode (+ve)
12	Opposite charges attract and like charges repel.
13	Products of salt solutions are harder to predict aswater also ionises to a very small extent producinghydrogen ions (H+) and hydroxide ions (OH-)
14	At the negative electrode (cathode): If there is morethan one type of positive ion in solution then theLEAST reactive element will be produced at the cathode.

lons	Cu2+(aq) and Cl1(aq) (from salt), H1(aq) and OH1(aq) (from water)		
cathode	Cu ² (aq) and H*(aq) ions are attracted. Copper ions are discharged more readily than hydrogen ions, so copper is formed as a brown solid.	Cu²+(aq) + Ze → Cu(s) (reduction)	
anode	Cl ⁻ (aq) and OH ⁻ (aq) ions are attracted. Chloride ions are discharged more readily than hydroxide ions, so chlorine is formed as a pale green gas.	\mathbb{R} 2Cl:(aq) \rightarrow Cl ₂ (g) + 2e (oxidation)	
Overall	ill The copper chloride decomposes but the water does not change CuCl _s (aq) → Cu(s) + Cl _s (g)		

	At the positive electrode (anode): oxygenis
	At the positive electrode (anode). Oxygenis
1	produced(from hydroxide ions), unlesshalideions (chloride,bromide or iodide ions) arepresent.Inthat case, thenegatively charged halide ionsloseelectrons and formthe corresponding halogen (chlorine,bromine or iodine).
	Thereactivity series is a list of metals inorder
2	of reactivity , the most reactive are at thetop.
	orreactivity, the most reactive are at thetop.
	The reactivity series is determined by
	howvigorouslyanelements reacts withother
	substances e.g. water andacid. Elements higher up the
3	reactivity series
	can displace those below them.
	•
	e.g. Zn + CuSO4 Cu + ZnSO4
	More reactive zinc replaced copper.

lons	Na*(aq) and Cl*(aq) (from salt), H*(aq) and OH*(aq) (from water)		
cathode	Na'(aq) and H'(aq) ions are attracted. Hydrogen ions are discharged more readily than sodium ions, so hydrogen gas is formed.	2H'(aq) + 2e → H ₂ (g) (reduction)	
anode	Cl (aq) and OH (aq) ions are attracted. Chloride ions are discharged more readily than hydroxide ions, so chlorine is formed as a pale green gas.	M 2Cl (aq) → Cl ₂ (g) + 2e (oxidation)	
Overall	The sodium chloride decomposes to form hydrogen and chlorine. The sodium and hydroxide ions remain in the solution. $2NaCl(aq) + 2H_2O(l) \rightarrow H_2(g) + Cl_2(g) + 2NaOH(aq)$		

4	Displacement reactions are also redox reactions. The reaction between zinc and copper sulphate above can be written as an ionic equation: Zn + Cu2+ Cu + Zn2+ The copper has gained two electrons this is red uction, zinc has lost two electrons to become Zn2+this is ox idation. This happens at the same time so a redox reaction.
5	Anore is a rock containing enough of a compound (or element) to be profitable when its sold. Very unreactive metals can be found their own or native state.e.g.gold and platinum.
6	The metals below carbon in the reactivity series e.g. iron are extractedby smelting. Metals above carbon e.g. aluminium by electrolysis. Both process use a lot energy and cause a lot of pollution. Biological methods of extractione.g. bioleaching and phytoextraction are less damaging to the environment and can work on ores with less metal in them.

Metal	Reaction with water	Reaction with dilute acid	Tendency of metal atoms to form cations	
potassium	react with cold water	react violently		
sodium	to form hydrogen and			
calcium	a metal hydroxide react to form		1 day	
magnesium	react very slowly, if	hydrogen	creasing ability of metal at form positive ions	
aluminium	at all, with cold water but react with steam	and a salt solution		
zinc	to form hydrogen and			
iron	a metal oxide			
copper	do not react with cold	do not react		
silver	water or steam			
gold			€ 9	



Most metals react with oxygen in the air, this is calledcorrosion. Paint, coating ofoiland asacrificial blockof a more reactive metal can stopcorrosion.An oxide layer (atarnish) also stops further reaction Corrosionofironneedswaterand is called rusting. A life cycle assessment (LCA) Is used to work out the environmentalimpact of a product. Recyclingcan make natural reserves of metal ore last longer, reduced mining of ores results in less damage to habitats, noise and dust.Less energy is needed so less carbon dioxide (global warming) and sulphur dioxide (acid rain) is emitted.

